arXiv:0904.0073v1 [math.GN] 1 Apr 2009

Stone-Čech compactifications and homeomorphisms of products of the long line

Veerendra Vikram Awasthi and Parameswaran Sankaran
Institute of Mathematical Sciences
CIT Campus, Chennai 600 113, INDIA
E-mail: vvawasthi@imsc.res.in
sankaran@imsc.res.in

Abstract: In this note we shall prove that the Stone-Čech compactification of $\mathcal{L}^n$ is the space $\overline{\mathcal{L}}^n$ where $\overline{\mathcal{L}}$ is the extended long line, namely, $\mathcal{L}$ together with its ends $\pm\Omega$. We give a similar description for the Stone-Čech compactification of the cartesian power of the semi-closed half-long line $\mathcal{L}_+$. As an application we show that any torsion subgroup of the group of all homeomorphisms of $\mathcal{L}_+^n$ (resp. $\mathcal{L}^n$) is isomorphic to a subgroup of the symmetric group S_n (resp. the semidirect product $(\mathbb{Z}/2\mathbb{Z})^n \ltimes S_n$).

1 Introduction

The well-known classification theorem for one-dimensional manifolds is that there are exactly two non-metrizable manifolds of dimension 1 (besides the two metrizable ones $\mathbb{R}$ and $\mathbb{S}^1$) namely the Alexandrof's long line and the half-long line. The half-long line is described as the space $\mathcal{L}_+ \setminus \{0\}$ where $\mathcal{L}_+ := [0,1) \times S_\Omega$ with lexicographic order topology and $0 := \{(0,0)\}$ is the smallest element. Here the S_Ω, the set of all ordinals which are less than Ω, is given the order topology, Ω being the smallest uncountable ordinal. The long line is then the space $\mathcal{L} = \mathcal{L}_- \cup \mathcal{L}_+$ glued at $\{0\}$ where $\mathcal{L}_-$ stands for $\mathcal{L}_+$ with its order reversed. The topology of these spaces have been studied mainly for their pathological properties in contrast with the well-behaved metrizable manifolds. However, we shall see that the Stone-Čech compactifications of $\mathcal{L}_+^n$ and $\mathcal{L}^n$ have elegant descriptions. Our main application is to problem of the characterizing, up to isomorphism, the torsion subgroups of the group of all homeomorphisms of $\mathcal{L}_+^n$ and $\mathcal{L}^n$.

Recall that the Stone-Čech compactification of $\mathcal{L}_+$ equals its one-point compactification, which is the extended half-long line $\overline{\mathcal{L}}_+ := \mathcal{L}_+ \cup \{\Omega\}$, and that of $\mathcal{L}$ is the extended long line $\overline{\mathcal{L}} := \mathcal{L}_+ \cup \{\Omega, -\Omega\}$, which is also its Freudenthal's end-compactification.

Theorem 1.1. *The Stone-Čech compactifications of $\mathcal{L}_+^n$ and $\mathcal{L}^n$ are respectively $\overline{\mathcal{L}}_+^n$ and $\overline{\mathcal{L}}^n$ for $n \geq 1$.*

AMS Mathematics Subject Classification (2000): 54D35, 55M35.
Keywords: Long line, Stone-Čech Compactification, homeomorphism groups

Although this is given as an exercise in [4, p.255-256] when $n = 1, 2$, we could not find in the literature the result stated for a general n.

We denote by Homeo(X) the group of all homeomorphisms of X. The symmetric group S_n acts on $(\mathbb{Z}/2\mathbb{Z})^n$ by permuting the factors. We denote by G_n the semi-direct product $(\mathbb{Z}/2\mathbb{Z})^n \ltimes S_n$. Note that S_n acts on $\mathcal{L}_+^n$ and $\mathcal{L}^n$ by permuting the coordinates. Also there is an obvious involution on $\mathcal{L}$ which yields an action of $(\mathbb{Z}/2\mathbb{Z})^n$ on $\mathcal{L}^n$, $n \geq 1$. This, together with the action of S_n, defines an action of G_n on $\mathcal{L}^n$.

Our next result is:

Theorem 1.2. *There exist surjective homomorphisms* Φ: Homeo$(\mathcal{L}_+^n) \longrightarrow S_n$ *and* Ψ: Homeo$(\mathcal{L}^n) \longrightarrow G_n$ *which split. Furthermore,* $\ker(\Phi)$ *and* $\ker(\Psi)$ *are torsion-free.*

An immediate corollary is the following:

Theorem 1.3. *Let* T *be a subgroup of* Homeo$(\mathcal{L}^n)$ *(resp.* Homeo$(\mathcal{L}_+^n)$ *in which every element is of finite order. Then* T *is isomorphic to a subgroup of* G_n *(resp.* S_n*).* □

Our proofs involve mostly elementary concepts from set-topology and make repeated use of the property that $\mathcal{L}_+$ and $\mathcal{L}$ are sequentially compact. We shall also make repeated use of [6, Theorem 1, §5.5] in §3.

For values of $n \leq 3$ (resp. $n \leq 2$), order of torsion elements of Homeo$(\mathcal{L}_+^n)$ (resp. $\mathcal{L}^n$)) have been determined by Deo and Gauld [3]. After this paper was completed, Satya Deo pointed out to us the preprint [2] in which the orders of torsion elements of Homeo$(\mathcal{L}_+^n)$ and Homeo$(\mathcal{L}^n)$, $n \geq 1$, have been determined, but not the *structure* of torsion subgroups.

2 Proof of Theorem 1.1

We use the following notations throughout. As usual I denotes the interval $[0, 1] \subset \mathbb{R}$. If M is a manifold with boundary (which could be empty), ∂M will denote the boundary of M. If $x \in \mathcal{L}$, then $-x \in \mathcal{L}$ denotes the image of x under the order reversing involution $\mathcal{L} \longrightarrow \mathcal{L}$ which switches $\mathcal{L}_+$ and $\mathcal{L}_-$ fixing 0. We denote by $\delta\overline{\mathcal{L}}_+^n$ the space $\overline{\mathcal{L}}_+^n \setminus \mathcal{L}_+^n$. Let $x = (x_1, \cdots, x_n) \in \overline{\mathcal{L}}_+^n$. $\Omega(x)$ denotes the set $\{j | x_j = \Omega\} \subset \{1, 2, \cdots, n\}$ and Ω_n denotes the point $(\Omega, \cdots, \Omega) \in \overline{\mathcal{L}}_+^n$. For any set-map $h : X \longrightarrow I$, and any subset $J \subset \mathbb{R}$, $h^{-1}(J)$ will have the obvious meaning, namely, $h^{-1}(I \cap J)$.

Let $x \in \delta\overline{\mathcal{L}}_+^n$. There is an obvious embedding $\gamma_x \colon \mathcal{L}_+ \longrightarrow \mathcal{L}_+^n$ defined as $t \mapsto x(t)$ where $x_i(t) = t$ if $i \in \Omega(x)$ and $x_i(t) = x_i$ if $i \notin \Omega(x)$. Let $f : \mathcal{L}_+^n \longrightarrow I$ be any continuous function. Since $f \circ \gamma_x : \mathcal{L}_+ \longrightarrow I$ is eventually constant, it can be extended continuously to $\overline{\mathcal{L}}_+$; its value at Ω will be denoted $\lim_{t \to \Omega} f(\gamma_x(t))$. Define $\widetilde{f} \colon \overline{\mathcal{L}}_+^n \longrightarrow I$ to be the set-map which extends f where $\widetilde{f}(x) := \lim_{t \to \Omega} f(\gamma_x(t))$ for all $x \in \delta\overline{\mathcal{L}}_+^n$. In the case of $\mathcal{L}_+^n$, the proof of Theorem 1.1 involves showing that $\widetilde{f}$ is continuous. The proof will be preceded by a couple of observations.

Lemma 2.1. *Let* $n \geq 1$ *and let* $h : \overline{\mathcal{L}}_+^n \longrightarrow I$ *be any continuous map. Then* h *is constant in a neighbourhood of* $\Omega_n = (\Omega, \ldots, \Omega)$*. Also any continuous map* $g : \overline{\mathcal{L}}^n \longrightarrow I$ *is constant in a neighbourhood of* $\{-\Omega, \Omega\}^n$*.*

Proof. Let $h(\Omega_n) = t_0$. Choose $\lambda_k < \Omega$ such that the basic open set $U_k = (\lambda_k, \Omega]^n$ is contained in $h^{-1}((t_0 - 2^{-k}, t_0 + 2^{-k}))$, $k \geq 1$. Let λ_0 be the least upper bound of $\{\lambda_k \mid k \geq 1\}$. If $p \in U_{\lambda_0}$, then $|h(p) - t_0| \leq 2^{-k}$ for every $k \geq 1$. Thus $h(p) = h(\Omega_n)$. Proof of the assertion concerning g is entirely similar and hence omitted. □

Lemma 2.2. *Let $n \geq 1$ and let $f : \mathcal{L}^n_+ \longrightarrow I$ be any continuous map. Then $\widetilde{f}$ defined above is continuous at $\Omega_n \in \overline{\mathcal{L}}^n_+$.*

Proof. Let $t_0 = \widetilde{f}(\Omega_n) = \lim_{\lambda \to \Omega} f(\lambda, \ldots, \lambda)$. First we shall show that given any $\epsilon > 0$, there exists a $\mu < \Omega$ such that $(\mu, \Omega)^n \subset \widetilde{f}^{-1}((t_0 - \epsilon, t_0 + \epsilon))$. Indeed we shall assume that there is no such μ and arrive at a contradiction.

Choose $\lambda_0 < \Omega$ such that $f(\lambda, \ldots, \lambda) = t_0$ for $\lambda_0 < \lambda < \Omega$. Assume that there exists an $\epsilon > 0$ such that $\widetilde{f}^{-1}((t_0 - \epsilon, t_0 + \epsilon))$ does contain $(\lambda, \Omega)^n$ for any $\lambda < \Omega$. Choose a $p_1 = (p_{1,1}, \ldots, p_{1,n}) \in (\lambda_0, \Omega)^n$ such that $|f(p_1) - t_0| \geq \epsilon$. Set $\lambda_1 := \max\{p_{1,j} \mid 1 \leq j \leq n\}$ and choose $p_2 = (p_{2,1}, \ldots, p_{2,n})$ such that $\lambda_1 < p_{2,j} < \Omega \; \forall j$ and $|f(p_2) - t_0| \geq \epsilon$. Continuing thus, we obtain a sequence $\lambda_0 < \lambda_1 < \cdots < \Omega$ and a sequence of points $p_1, p_2, \cdots$ in $\mathcal{L}^n_+$ such that $\lambda_k = \max\{p_{kj} \mid 1 \leq j \leq n\}, \lambda_k < p_{k+1,j}, 1 \leq j \leq n$, and $|f(p_j) - t_0| \geq \epsilon$ for all $j \geq 1$. It is evident that the sequence (p_k) has limit $p = (\nu, \ldots, \nu) \in \mathcal{L}^n_+$ where $\nu = \lim \lambda_k$. Hence by continuity of f at p, we obtain $|f(p) - t_0| \geq \epsilon$. But this is a contradiction since, by our choice of λ_0, we have $f(\nu) = t_0$ as $\nu > \lambda_0$. Hence $\widetilde{f}^{-1}((t_0 - \epsilon, t_0 + \epsilon))$ must contain $(\mu, \Omega)^n$ for some $\mu < \Omega$ as claimed. Now choose such a μ.

We claim that $(\mu, \Omega]^n \subset \widetilde{f}^{-1}((t_0 - \epsilon, t_0 + \epsilon))$. To see this observe that if $x \in (\mu, \Omega]^n$ with $\Omega(x) \neq \emptyset$, then $\gamma_x(t) \in (\mu, \Omega)^n$ for all $t > \mu$. Therefore $|t_0 - f(\gamma_x(t))| < \epsilon$ for $t > \mu$. Since $f \circ \gamma_x$ is eventually constant, we have $\widetilde{f}(x) = \lim_{t \to \Omega} f(\gamma_x(t)) = f(\gamma_x(\lambda))$ for some $\lambda > \mu$ and so $|\widetilde{f}(x) - t_0| < \epsilon$. This completes the proof. □

Proof of Theorem 1.1: We prove the theorem by induction on n. As remarked already, it is a well-known property when $n = 1$. We shall only consider the case of $\mathcal{L}^n_+$. The proof in the case of $\mathcal{L}^n$ involves only obvious and routine changes.

Now suppose that $n \geq 2$ and that the theorem holds for all positive integers up to $n - 1$. Let $p = (p_1, \cdots, p_n) \in \delta\overline{\mathcal{L}}^n_+$. Then $\Omega(p) = \{j \mid p_j = \Omega\} \neq \emptyset$. Continuity at $p = \Omega_n$ having been established in Lemma 2.2, we assume that $1 \leq k := n - \#\Omega(p) < n$.



Figure 1: $J \times \overline{\mathcal{L}}^{n-k}_+$.

For convenience we shall suppose that $\Omega(p) = \{j \mid k+1 \leq j \leq n\}$. Set $a := (p_1, \ldots, p_k) \in \mathcal{L}_+^k$. Choose a 'small' basic open set $J \subset \mathcal{L}_+^k$ containing a, i.e., J is a product of bounded open intervals in $\mathcal{L}_+$. Let A be a countable dense subset of J such that $a \in A$. For any $b \in J$, let f_b denote the restriction of f to the slice $Y_b := \{b\} \times \mathcal{L}_+^{n-k}$. By inductive hypothesis $\widetilde{f_b} : \{b\} \times \overline{\mathcal{L}}_+^{n-k} \longrightarrow I$ is continuous. Now by Lemma 2.1 $\widetilde{f_b}$ is constant in a neighbourhood of (b, Ω_{n-k}). For each $b \in A$, choose an element $\lambda_b < \Omega$ such that $\widetilde{f_b}$ is constant on $\{b\} \times U_{\lambda_b}$ where $U_\lambda = \{q \in \overline{\mathcal{L}}_+^{n-k} \mid q_i > \lambda \ \forall i\}$. Since the $\{\lambda_b \mid b \in A\}$ is a bounded countable set, it has a least upper bound $\mu < \Omega$. Now $\widetilde{f}$ restricted to $A \times U_\mu$ factors through the projection $A \times U_\mu \longrightarrow A$. That is, $\widetilde{f}(b, x) = f(b, \mu)$ for all $b \in A$ and $x \in [\mu, \Omega]$. Since $A \times U_\mu$ is dense in $J \times U_\mu$, this last equality holds for all $b \in J$ and $x \geq \mu$ by continuity of f. It follows that $\widetilde{f}$ is continuous at p. □

We remark that by our proof of Theorem 1.1 and Lemma 2.1, *any* continuous map $\mathcal{L}_+^n \longrightarrow I$ is constant in a basic open set $(\mu, \Omega)^n$ for some μ. In fact, as a corollary of the proof of Theorem 1.1, we obtain the following proposition. Let $p = (\alpha, \cdots, \alpha) \in \mathcal{L}_+^n$ be any point on the diagonal. Then there is a retraction $r_\alpha : \overline{\mathcal{L}}_+^n \longrightarrow I_\alpha^n$ where $I_\alpha = [0, \alpha]^n \cong I^n$. Specifically, let r_α be the product of retractions $\overline{\mathcal{L}}_+ \longrightarrow I_\alpha$ which sends any $\beta \geq \alpha$ to α. Similarly, one has the retraction $\overline{\mathcal{L}} \longrightarrow J_\alpha := [-\alpha, \alpha] \cong I$ defined by $\beta \mapsto \alpha$ and $-\beta \mapsto -\alpha$ if $\beta \geq \alpha$. This defines a retraction $\rho_\alpha : \overline{\mathcal{L}}^n \longrightarrow J_\alpha^n$.

Proposition 2.3. *Any continuous map $f : \overline{\mathcal{L}}_+^n \longrightarrow I$ factors through the retraction r_α for some $\alpha < \Omega$. Similarly, any continuous map $g : \overline{\mathcal{L}}^n \longrightarrow J_\alpha$ factors through the retraction ρ_α for some $\alpha < \Omega$.*

Proof. We shall prove the statement for f, the case of g being analogous.

Let $p \in \delta\overline{\mathcal{L}}_+^n = \overline{\mathcal{L}}_+^n \setminus \mathcal{L}_+^n$. If $p = \Omega_n$, let $\mu \in \mathcal{L}_+$ be such that $f|(\mu, \Omega]^n$ is constant.

Now let $p \neq \Omega_n$ so that $0 < \#\Omega(p) < n$. In the course of our proof of Theorem 1.1, it was observed that there is a basic open set $U(p) := \prod U_i(p)$ containing p where $U_i(p)$ is a bounded interval in $\mathcal{L}_+$ if $i \notin \Omega(p)$ and $U_i(p) = (\mu(p), \Omega]$ if $i \in \Omega(p)$ such that $f|U(p)$ factors through the projection $U(p) \longrightarrow \prod_{j \notin \Omega(p)} U_j(p) \prod_{j \in \Omega(p)} \{\mu(p)\}$.

The collection $\{U(p)\}, p \in \delta\overline{\mathcal{L}}_+^n$, is an open covering of the compact space $\delta\overline{\mathcal{L}}_+^n$. Therefore there exists finitely many points $p_1, \cdots, p_k$ such that $\{U(p_j)\}$ covers $\delta\overline{\mathcal{L}}^n$. Now let $\mu = \max\{\mu(p_j) \mid 1 \leq j \leq k\}$. It is straightforward to verify that f factors through r_μ. □

Remark 2.4. The same arguments as in the proof of Theorem 1.1 also shows that the Stone-Čech compactification of $\mathcal{L}_+^n \times I$ is $\overline{\mathcal{L}}_+^n \times I$ and that of $\mathcal{L}^n \times I$ is $\overline{\mathcal{L}}^n \times I$. It follows readily that if $f, g : \mathcal{L}_+^n \longrightarrow \mathcal{L}_+^n$ are continuous, then f is homotopic to g if and only if their extensions $\widetilde{f}, \widetilde{g} : \overline{\mathcal{L}}_+^n \longrightarrow \overline{\mathcal{L}}_+^n$ are. An analogous statement holds for self-maps of $\mathcal{L}^n$.

3 Homeomorphisms of $\mathcal{L}_+^n$ and $\mathcal{L}^n$

Observe that $\overline{\mathcal{L}}_+^n, \overline{\mathcal{L}}^n, n \geq 1$, are not path connected. The path components of $\overline{\mathcal{L}}_+^n$ are labelled by the set $V_n := \{0, \Omega\}^n$. More precisely, the elements of V_n are in distinct path components and every

path component of $\overline{\mathcal{L}}_+^n$ contains a point of V_n. We shall denote the path component containing $p \in V_n$ by X_p. Note that $X_p = \prod U_j$ where $U_j = \mathcal{L}_+$ if $j \notin \Omega(p)$ and $U_j = \{\Omega\}$ otherwise. In particular, $\dim X_p = n - \#\Omega(p)$.

We obtain a directed graph $\mathcal{H}_n$ (or just $\mathcal{H}$) whose vertex set is $\mathcal{H}^0 = \{X_p \mid p \in V_n\}$ and edge set $\mathcal{H}^1 = \{\epsilon(p,q) \mid X_p \subset \overline{X}_q, \dim X_p = \dim X_q - 1\}$. The edge $\epsilon(p,q)$ is oriented so that it issues from X_p to X_q. One has a partition of $V_n = \cup_{0\le k\le n} V_n(k)$ where $V_n(k) = \{p \in V_n \mid \#\Omega(p) = k\}$. The number of edges issuing from (resp. terminating at) X_p equals $\#\Omega(p)$ (resp. $\dim X_p$). We let H_n denote the group of all automorphisms of the directed graph $\mathcal{H}_n$. Every element of H_n fixes $X_{\Omega_n} = \{\Omega_n\}$. It is not hard to see that H_n is isomorphic, via restriction, to the group of permutations of $\{X_p \mid p \in V_n(n-1)\} \subset \mathcal{H}_n^0$. Thus $H_n \cong S_n$.

Let $h : \overline{\mathcal{L}}_+^n \longrightarrow \overline{\mathcal{L}}_+^n$ be a homeomorphism and let h_* denote the induced map on the set of path-components of $\overline{\mathcal{L}}_+^n$.

Proposition 3.1. *Any homeomorphism $h: \overline{\mathcal{L}}_+^n \longrightarrow \overline{\mathcal{L}}_+^n$ induces an isomorphism of the directed graph $\mathcal{H}_n$. The map $h \mapsto h_*$ is a surjective homomorphism of groups $\Phi : Homeo(\overline{\mathcal{L}}_+^n) \longrightarrow H_n$ which splits.*

Proof. Any homeomorphism of $\overline{\mathcal{L}}_+^n$ induces an isomorphism of the set of path components. Furthermore this defines a homomorphism from the group Homeo$(\overline{\mathcal{L}}_+^n)$ to the group of permutations of the set of path-components of $\overline{\mathcal{L}}_+^n$. So $h \in$Homeo$(\overline{\mathcal{L}}_+^n)$ induces a bijection of the vertex set the graph $\mathcal{H}_n$. Also $h(\overline{X}_p) \subset h(X_q)$ if $\overline{X}_p \subset X_q$. It follows that it preserves the oriented edges of $\mathcal{H}_n$. Hence h induces an isomorphism h_* of $\mathcal{H}_n$.

Clearly every homeomorphism of $\overline{\mathcal{L}}_+^n$ fixes Ω_n. Hence h maps $V_n(n-1)$ onto $V_n(n-1)$. Any permutation σ of $V_n(n-1)$ is evidently realizable by the homeomorphism h of $\overline{\mathcal{L}}_+^n$ given by the *same* permutation of the coordinates. Therefore Φ is surjective. This also shows that Φ splits. □

Now consider the space $\overline{\mathcal{L}}^n$. Since the path components of $\overline{\mathcal{L}}$ are $-\Omega, \Omega$ and $\mathcal{L}$, $\overline{\mathcal{L}}^n$ has 3^n path components. They are labelled by $\{-\Omega, 0, \Omega\}^n$. The element $q \in \{-\Omega, 0, \Omega\}^n$ labels the path component $X_q = \prod U_j$ where $U_j = \{q_j\}$ if $q_j \neq 0$ and $U_j = \mathcal{L}$ if $q_j = 0$. Observe that $\dim X_q = \#\{j \mid \dim q_j = 0\}$.

Let $W_n = \{-\Omega, \Omega\}^n$. Each element of W_n forms a path component of $\overline{\mathcal{L}}^n$. Observe that *any* self-homeomorphism h of $\overline{\mathcal{L}}^n$ preserves W_n as the other path components are of positive dimension.

Consider the (simple) graph $\mathcal{G}_n$ whose vertices are X_p, $p \in W_n$. The edges of the graph are $e_{p,q}$ if p and q differ exactly in one coordinate (where they differ by sign). The group of automorphisms of $\mathcal{G}_n$ is isomorphic to the semi-direct product $(\mathbb{Z}/2\mathbb{Z})^n \ltimes S_n := G_n$ where the actions of $(\mathbb{Z}/2\mathbb{Z})^n$ and S_n are obtained from their obvious respective actions on W_n.

Proposition 3.2. *Any homeomorphism h of $\overline{\mathcal{L}}^n$ induces an isomorphism h_* of the graph $\mathcal{G}_n$. Furthermore, $h \mapsto h_*$ defines a surjective homomorphism of groups $\Psi : Homeo(\overline{\mathcal{L}}^n) \longrightarrow G_n$ which splits.*

Proof. As observed above, $h(W_n) = W_n$ and so h induces a bijection of the vertices of $\mathcal{G}_n$. Suppose that $e_{p,q}$ is an edge of $\mathcal{G}_n$, say, $p_i = \Omega = -q_i, p_j = q_j$ for $j \neq i$. Consider $X_{p,q} = \{x \in \overline{\mathcal{L}}^n \mid x_i \in \mathcal{L}, x_j = p_j, j \neq i\} \cong \mathcal{L}$. Then $h(X_{p,q})$ has to be a path component of dimension 1 which contains

$h(p), h(q) \in W_n$ in its closure. It follows that $h(p), h(q)$ are end points of an edge of $\mathcal{G}_n$. Therefore h_* is an isomorphism of $\mathcal{G}_n$.

It is evident that the homeomorphisms of $\overline{\mathcal{L}}^n$ which flips the signs of certain coordinates forms a subgroup of Homeo($\mathcal{L}^n$) isomorphic to $(\mathbb{Z}/2\mathbb{Z})^n$. These, together with the homeomorphisms which permute the coordinates form a group isomorphic to $(\mathbb{Z}/2\mathbb{Z})^n \ltimes S_n$. It is evident that Ψ maps this subgroup isomorphically onto G_n. Therefore Ψ is split. □

Lemma 3.3. *Let M be a connected manifold with non-empty boundary ∂M. Suppose that $f\colon M \longrightarrow M$ is a homeomorphism which restricts to the identity on ∂M. Then either f is the identity or is of infinite order.*

Proof. Consider the double $\hat{M} = M_0 \cup_{\partial M} M_1$ of M, obtained by gluing two copies M_0, M_1 of M along the common boundary. Since $f|\partial M$ is identity, it extends to a homeomorphism $f_0 : \hat{M} \longrightarrow \hat{M}$ where f_0 is just f on M_0 and is identity on M_1. Since f_0 is identity on a non-empty open set of $\hat{M}$, it follows that that f_0 is of infinite order (cf. [6, Theorem 1, §5.5]). Hence f has to be of infinite order. □

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2: In view of Theorem 1.1 any homeomorphism of $\mathcal{L}^n_+$ lifts to a unique homeomorphism of $\overline{\mathcal{L}}^n_+$ and similarly any homeomorphism of $\mathcal{L}^n$ lifts to a unique homeomorphism of $\overline{\mathcal{L}}^n$. Indeed one has, in fact, isomorphisms of groups Homeo($\mathcal{L}^n_+$) ≅Homeo($\overline{\mathcal{L}}^n_+$) and Homeo($\mathcal{L}^n$) ≅Homeo($\overline{\mathcal{L}}^n$).

By Proposition 3.1 one has a surjective homomorphisms of groups Homeo($\overline{\mathcal{L}}^n_+$)$\longrightarrow H_n \cong S_n$, which splits, defined as $h \mapsto h_*$, where H_n is the group of automorphisms of the directed graph $\mathcal{H}_n$. Similarly Homeo($\overline{\mathcal{L}}^n$)$\longrightarrow G_n$ defined by $g \mapsto g_*$ is a surjective homomorphism of groups by Proposition 3.2. As observed already, $H_n \cong S_n$ and $G_n \cong (\mathbb{Z}/2\mathbb{Z})^n \ltimes S_n$.

To complete the proof, we need only to show that ker(Φ) and ker(Ψ) are torsion-free. When $n = 1$ the statement is trivial to verify. Assume that $n > 1$ and that the theorem is valid for all dimensions up to $n - 1$.

Let f be any element of Homeo($\overline{\mathcal{L}}^n_+$) of finite order such that $f_* \in H_n$ is trivial. Observe that $X_0 = \mathcal{L}^n_+$. We shall show that $f|\partial X_0$ is the identity homeomorphism. It would then follow, in view of Lemma 3.3, that $f|X_0$ is identity and so f itself is identity as X_0 is dense in $\overline{\mathcal{L}}^n_+$.



Figure 2: ∂X_0.

Note that f maps each path component of $\overline{\mathcal{L}}^n_+$ to itself and $f_p := f|X_p$ is a finite order homeomorphism of X_p for each $p \in V_n$. Let $k = \#\Omega(p)$. If $k > 0$, then $X_p \cong \mathcal{L}^{n-k}_+$ and furthermore f_p induces the identity map of the directed graph $\mathcal{H}_{n-k}$ associated to X_p. Hence, by induction hypothesis, f_p is the identity map of X_p. Now let $k = 0$. In this case $p = 0$ and ∂X_0 is *not* homeomorphic to $\mathcal{L}^n_+$ and so we cannot apply inductive hypothesis directly to conclude that $f|\partial X_0$ is the identity map. However, note that the points $p \in V_n(n-1)$ in the closure $\overline{\partial X}_0$ of ∂X_0 have basic neighbourhoods in $\overline{\partial X}_0$ whose closures are homeomorphic to $\overline{\mathcal{L}}^{n-1}_+$. Consider, say, the point $p = (\Omega, \cdots, \Omega, 0) \in \overline{\partial X}_0$. By what has been shown already, we have $f(p) = p$. Since f has finite order, there exists a neighbouhood $U_0 \subset \overline{\partial X}_0$ of p which is invariant under f. Choose a $\lambda_0 < \Omega$ such that the basic open set $B_0 := (\lambda_0, \Omega]^{n-1} \times \{0\}$ is contained in U_0. The open set $\cap_{0 \le j < r} f^r(B_0) =: U_1$ is invariant under f where r is the order of f. Repeating this argument, we get a sequence $\lambda_0 < \lambda_1 < \cdots < \Omega$ in $\mathcal{L}_+$ and open sets $U_0 \supset B_0 \supset U_1 \supset B_1 \supset \cdots$ in $\overline{\partial X}_0$ where $B_i = (\lambda_i, \Omega]^{n-1} \times \{0\}$ and U_i are invariant under f. Let $\mu < \Omega$ be the limit of (λ_i). Then $C := \cap_{k \ge 0} U_k = \cap_{k \ge 0} B_k = [\mu, \Omega]^{n-1} \times \{0\}$ is invariant under f and $C \cong \overline{\mathcal{L}}^{n-1}_+$. Note that $(f|C)_*$ is the identity automorphism of the directed graph $\mathcal{H}_{n-1}$ associated to C since $f(x) = x$ for all $x \in X_q \cap C$ for all $q \in V_n(k), k \ge 1$. Since $f|C$ is of finite order, by induction hypothesis we conclude that $f|C$ is trivial. Now by [6, Theorem 1, §5.5] we conclude that f is the identity map.

Proof in the case of $\mathcal{L}^n$ is similar and we merely give an outline. Let g be a finite order homeomorphism of $\overline{\mathcal{L}}^n$ which induces the identity automorphism of $\mathcal{G}_n$. Let X be any path component of $\overline{\mathcal{L}}^n$ which is of dimension less than n. If X is zero-dimensional, then it is point-wise fixed by g as g_* is the identity. Otherwise. $X \cong \mathcal{L}^k, 1 \le k < n$, the map $g|X$ is of finite order and induces the identity map of the graph associated to X. Hence, by induction hypothesis, $g|X$ is identity. Thus $g|(\delta\overline{\mathcal{L}}^n)$ is the identity. Now consider $\Omega_n \in \overline{\mathcal{L}}^n$. Proceeding as in the construction of C above, we obtain a $\mu < \Omega$ such that $g(D) = D$ where $D := [\mu, \Omega]^n \subset \overline{\mathcal{L}}^n$. Note that $D \cong \overline{\mathcal{L}}^n_+$. Now $g|D$ is a finite order element and it induces identity map of the directed graph $\mathcal{H}_n$. Hence, by what has been shown already, $g|D$ is identity. Now by [6, Theorem 1, §5.5], $g|\mathcal{L}^n$ is identity. □

Remark 3.4. (i) Let $h : \overline{\mathcal{L}}^n_+ \longrightarrow \overline{\mathcal{L}}^n_+$ be a homeomorphism. The induced automorphism h_* of the directed graph $\mathcal{H}_n$ determines and is determined by the isomorphism $H_0(h) : H_0(\overline{\mathcal{L}}^n_+; \mathbb{Z}) \longrightarrow H_0(\overline{\mathcal{L}}^n_+; \mathbb{Z})$ induced by h in 0-th singular homology. Since $H_0(\overline{\mathcal{L}}^n_+)$ is the free abelian group on the set of vertices of $\mathcal{H}_n$, we see that the elements of $S_n \subset \mathrm{Homeo}(\overline{\mathcal{L}}^n_+)$ are in distinct homotopy classes. In view of Remark 2.4, we see that distinct elements of $S_n \subset \mathrm{Homeo}(\mathcal{L}^n_+) \cong \mathrm{Homeo}(\overline{\mathcal{L}}^n_+)$ are in distinct homotopy classes. The last statement also holds for the group $G_n \subset \mathrm{Homeo}(\mathcal{L}^n)$ and can be seen by a similar argument. See also [1, p. 44] and [2]. It follows that the homomorphisms Φ and Ψ factor through the mapping class groups of $\mathcal{L}^n_+$ and $\mathcal{L}^n$ respectively. It is shown in [2] that mapping class groups of $\mathcal{L}^n_+$ and $\mathcal{L}^n$ are isomorphic to H_n and G_n respectively. Thus $\ker(\Phi)$ and $\ker(\Psi)$ consist precisely of homeomorphisms which isotopic to respective identity maps.

(ii) It is an interesting problem to classify *conjugacy classes* of finite subgroups of $\mathrm{Homeo}(\mathcal{L}^n)$ and of $\mathrm{Homeo}(\mathcal{L}^n_+)$.

Acknowledgements: The authors thank Satya Deo for a talk he gave at the Ramanujan Institute for Advanced Study in Mathematics, Chennai, in March 2008, which initiated our interest in homeomorphism groups of non-metrizable manifolds. We thank him also for providing us a copy of [3]. The second author gratefully acknowledges financial support from Abdus Salam International

Centre for Theoretical Physics, Trieste, Italy, during his visit in the autumn of 2008, where part of this work was done.

References

[1] M. Baillif, The homotopy classes of continuous maps between some nonmetrizable manifolds. Topology Appl. 148 (2005), no. 1-3, 39–53.

[2] M. Baillif, S. Deo, and D. Gauld, The mapping class group of the powers of the long ray and other non-metrizable spaces, unpublished, 2006.

[3] Satya Deo and David Gauld, The torsion of the group of homeomorphisms of powers of the long line. J. Aust. Math. Soc. **70** (2001), 311–322.

[4] J. Dugundji, *Topology*, Allyn and Bacon, Inc., Boston, Mass. 1966.

[5] David Gauld, Homeomorphisms of 1-manifolds and ω-bounded 2-manifolds. *Papers on general topology and applications.* (Madison, WI, 1991), 142–149, Ann. New York Acad. Sci., **704**, 1993.

[6] D. Montgomery and L. Zippin, *Topological transformation groups*, Interscience, 1955.